



The outcome of pregnancy in a kidney transplant patient: a case report and review of the literature

Ishod trudnoće kod bolesnice sa transplantiranim bubregom: prikaz slučaja i pregled literature

Andreja Glišić*, Nevena Divac[†], Miroslava Gojnić Dugalić*,
Biljana Kastratović Kotlica*, Neven Vavić[‡], Nataša Cerovac[§], Milica Prostran[†]

University of Belgrade, Faculty of Medicine, Clinical Center of Serbia, *Clinic for Gynecology and Obstetrics, [†]Institute of Pharmacology, Clinical Pharmacology and Toxicology, Belgrade, Serbia; Military Medical Academy, [‡]Center for Transplantation of Solid Organs, Belgrade, Serbia; [§]Clinic for Neurology and Psychiatry for Children and Youth, Belgrade, Serbia

Abstract

Introduction. The possibility of a term pregnancy with favorable maternal and neonatal outcome is one of the greatest advances in kidney transplantation, though concerns still exist regarding the safety of the mother, fetus, and graft. The use of immunosuppressive medications during pregnancy is related to possible fetal adverse effects. **Case report.** We report a course of a pregnancy in a patient with a kidney transplant. The patient was treated with immunosuppressive therapy (tacrolimus, azathioprine, and prednisolone) during the pregnancy. The outcome of the pregnancy was without maternal and neonatal complications. Serum creatinine levels were stable and no acute organ rejection occurred during pregnancy. Significant elevation of the D-dimer and coagulant factors II, VII, IX and X were noticed

during the third trimester. This could be partially attributed to azathioprine, which was a part of the immunosuppressive regimen. On the other hand, there were no radiological or clinical signs of thromboembolism, but low-molecular-weight heparin prophylaxis was immediately initiated. Cesarean section was performed at the 39th gestational week and a healthy female infant was delivered with a birth weight of 3,150 g and Apgar score 9. **Conclusion.** Pregnancies of kidney transplant recipients are high-risk and require a multidisciplinary approach. Careful clinical follow-up is a prerequisite for favorable outcome.

Key words:
kidney transplantation; pregnancy; fetal development; tacrolimus; azathioprine; prednisolone.

Apstrakt

Uvod. Mogućnost uspešne trudnoće kod žena sa presađenim bubregom smatra se jednim od najvećih uspeha ove vrste lečenja, ali nosi sa sobom i određene probleme u vezi sa bezbednošću majke, fetusa i presađenog organa. Upotreba immunosupresivne terapije tokom trudnoće povezana je sa mnogim neželjenim efektima. **Prikaz slučaja.** Prikazali smo bolesnicu sa presađenim bubregom koja je tokom trudnoće primala immunosupresivnu terapiju (takrolimus, azatioprin i prednizolon). Ishod trudnoće je bio uspešan i po majku i po novorođenče. Nivo serumskog kreatinina majke bio je stabilan sve vreme trudnoće i nije doslo do akutnog odbacivanja presađenog organa. Tokom trećeg trimestra

došlo je do neuobičajenog porasta D-dimera i faktora koagulacije II, VII, IX i X, što je retka, ali moguća komplikacija primene azatioprina. Nije bilo kliničkih niti radioloških znakova tromboembolizma, ali je niskomolekularni heparin uveden profilaktički. Trudnoća je završena u 39-oj nedelji gestacije planiranim carskim rezom i rođeno je zdravo žensko dete porođajne težine 3 150 g, ocenjeno Apgar skorom 9. **Zaključak.** Trudnoća kod bolesnice sa presađenim bubregom smatra se visoko rizičnom i zahteva pažljivo planiranje i praćenje.

Ključne reči:
transplantacija bubrega; trudnoća; trudnoća, razvoj fetusa; takrolimus; azatioprin; prednizolon.

Introduction

The impaired reproductive function is one of the detrimental consequences of end-stage renal disease (ESRD). The restoration of pituitary-ovarian function and fertility in female kidney transplant recipients is one of the greatest achievements of modern transplantation. The first report of a pregnancy in a kidney transplant recipient was published in 1958. After that, due to the improvement of immunosuppressive therapy and modern perinatal care, series of pregnancies with favorable outcomes have been reported worldwide¹⁻³. The pregnancy in a kidney transplant patient is complicated due to previous major abdominal surgery, pre-existing comorbidities such as diabetes, hypertension etc, and the adverse effects of immunosuppressive treatment³. Pregnancy complications associated with chronic renal failure, such as hypertension, proteinuria, preeclampsia and preterm delivery, are still possible in kidney transplant recipients. These pregnancies are always considered as high-risk and deserve careful planning and clinical follow-up from the very beginning.

Immunosuppressive medications prevent kidney rejection but also carry significant risks. During pregnancy, these medications are related to many serious fetal adverse effects. Therefore, it is important to carefully evaluate the safety and efficacy of immunosuppressive medication during pregnancy and to provide adequate clinical prenatal care in order to reduce the risk of graft rejection on one side, and unwanted effects on fetal development on the other side.

The present study is a review of a new case of a pregnancy in a kidney transplant recipient treated with tacrolimus, azathioprine, and prednisolone, with successful maternal and fetal outcome. There have been very few reports of such pregnancies in Serbia.

Case report

The patient was a 26 year old pregnant woman with the history of kidney disease. She was diagnosed with ESRD eight years ago. Hemodialysis was initiated three years later. After eight months of dialysis, she was transplanted a living related donor kidney at the Military Medical Academy in Belgrade. The procedure was performed successfully and the patient was discharged on the 23rd post-transplantation day with serum creatinine level of 161 $\mu\text{mol/L}$. The clinical protocol included immunosuppressive treatment comprising tacrolimus, prednisolone and mycophenolate mofetil. Antihypertensive therapy was prescribed (metoprolol and nifedipine) and on discharge, her blood pressure was within recommended values for kidney transplant patients⁴. The patient was followed-up at regular intervals and the doses of immunosuppressive drugs were gradually tapered to maintenance doses.

Three years later, the patients spontaneously conceived. She was referred to the Clinic of Gynaecology and Obstetrics, Clinical Center of Serbia at 5 weeks gestation. Since pregnancies in transplant patients require a multidisciplinary approach, a close cooperation with nephrologists from the

Military Medical Academy was established. Her immunosuppressive regimen was changed to tacrolimus 3.5 mg in the morning and 3.5 mg in the evening, prednisolone 7.5 mg/day and azathioprine 50 mg/day. Mycophenolate mofetil was stopped. Beta blocker was also withdrawn, verapamil was introduced and methyl dopa continued. The patient was followed on monthly basis throughout the entire pregnancy. Ultrasonographic examinations showed normal fetal growth and amniotic fluid volume was in the normal range. The laboratory values of serum glucose, electrolytes, serum urea and creatinine, proteins in 24-hour urine collection and protein to creatinine ratio were regularly controlled. The serum creatinine levels throughout the course of pregnancy were stable between 79–110 $\mu\text{mol/L}$ and the day before the caesarean section it was 116 $\mu\text{mol/L}$. The urine protein to creatinine ratio was stable in range 0.38–0.41. Blood pressure was well controlled with methyl dopa and verapamil.

The patient's coagulation status was also closely monitored. During her third trimester, D-dimer and coagulation factors (II, VII, VIII, IX and X) were elevated. D-dimer levels were between 3.13 to 4.38 mg/L. The left-sided leg edema also developed. The lower-extremity venous duplex ultrasound was performed, and she was found to have no signs of deep venous thrombosis (DVT) or superficial thrombophlebitis. However, it was decided that anticoagulant prophylaxis should be started and low molecular weight heparin (fraxiparine) was initiated. Asymptomatic bacteriuria was treated by ceftriaxone 2g /daily i.m. during ten days.

We performed planned caesarean delivery because the transplant kidney was located very low in the pelvis and probably could obstruct the labor.

A healthy female child was born at term by caesarean section. Birth weight was 3,150 g, Apgar score 9 at first minute of life and no congenital malformations were identified. Neurological and clinical status of the baby at birth was normal. During first months of life, the baby achieved age-appropriate developmental milestones. During the early postoperative period, the patient's serum creatinine level showed transient elevation up to 218 $\mu\text{mol/L}$ but reverted to baseline values. We performed intravenous hydration (3,000 mL of fluid *per* a day) during 7 days. There were no signs of acute kidney rejection. The patient was advised against breastfeeding and bromocriptine was administered to terminate lactation.

Discussion

Pregnancy in kidney transplant recipients is burdened with risks and requires careful follow-up. These risks include impaired renal function, graft rejection, spontaneous abortion, preterm delivery, low birth weight and fetal growth retardation⁵. Great concern in such pregnancy is the potential adverse effect of immunosuppressant drugs to the fetus.

The recommended immunosuppressive regimen in pregnant kidney transplant recipients usually comprises a calcineurin inhibitor, corticosteroid and azathioprine⁶. Mycophenolate mofetil and mammalian target of rapamycin inhibitors (mTOR) are associated with increased incidence of spontaneous abortions and congenital malformations in fetuses^{7, 8}.

Therefore, these two drugs are not recommended during pregnancy and should be discontinued before conception, or, if the pregnancy was not planned, immediately after^{7,9}.

Tacrolimus belongs to a class of calcineurin inhibitors, together with cyclosporine. It falls into pregnancy category C by the U.S. Food and Drug Administration (FDA). Tacrolimus is preferred over cyclosporine due to better efficacy regarding graft function and graft survival¹⁰. Additionally, compared to tacrolimus, cyclosporine in pregnant women is associated with higher incidence of hypertension¹¹. Fetal levels of cyclosporine are similar to maternal levels, whereas the umbilical cord concentration of tacrolimus is 19% of maternal unbound plasma concentration. The lower fetal concentration of tacrolimus is attributed to the active efflux of tacrolimus from the fetus toward the mother by placental P-glycoprotein activity^{12,13}. Tacrolimus has been reported to cause hyperkalemia and kidney impairment in neonates, as well as premature delivery and preeclampsia^{13,14}. Teratogenicity in children whose mothers were treated with tacrolimus during pregnancy is not significantly increased compared to general population^{15,16}.

Prednisone has been widely used in pregnancy for indications other than solid organ transplantations. It is listed as pregnancy category C by the FDA¹⁷. Fully developed placenta partially protects the fetus from prednisone exposure by its metabolic activity (enzyme 11-beta-hydroxylase)¹⁸. The maternal to cord blood ratio of prednisone is 8 : 1 to 10 : 1¹⁹. However, the risk of neonatal adrenal insufficiency cannot be ruled out even with low doses of prednisone and careful post-natal monitoring is required.

Azathioprine and its active metabolite, 6-mercaptopurine, are purine analogues which interfere with the synthesis of adenine and guanine ribonucleosides²⁰. Azathioprine has been labeled as pregnancy category D by FDA²¹. It crosses the placenta. However, the fetus cannot metabolize it to its active metabolite 6-mercaptopurine due to lack of liver enzymes²². Azathioprine can cause hematological disturbances and immunodeficiency in the fetus^{11,23}. The frequency of congenital abnormalities in infants of kidney transplant recipient mothers was between 0.0–11.8% in different case-series studies²⁰. The prospective, controlled cohort study by Goldstein et al.²⁴ which compared pregnancy outcome in women exposed to azathioprine (n = 189) to non-

exposed controls (n = 230) showed no difference in terms of teratogenicity. However, the azathioprine exposed group had a higher incidence of low birth weight and prematurity.

Additionally, increased thrombotic complications in kidney transplant recipients have been associated with the use of azathioprine²⁵. One of the postulated mechanisms for this is the stimulation the synthesis of coagulation factors induced by azathioprine, especially factors II and X²⁶. It could partially explain the elevation of coagulation factors in our patient.

In women with a renal transplant, the cesarean delivery rate approaches 50%²⁷. We performed planned caesarean delivery. Transplant kidney was located very low in pelvis near to bladder and a low segment of uterus. We made a medial abdominal incision and medial uterine incision to protect the transplanted kidney. Probably, the transplanted kidney could obstruct the labor.

Our case emphasizes the significance of a multidisciplinary approach to a pregnancy in a kidney transplant recipient. Due to excellent coordination between gynecologists, nephrologists, pediatricians and clinical pharmacologists, our patient had a successful pregnancy with favorable maternal, fetal and graft outcome. The graft function remained stable as well as blood pressure and other clinically significant parameters in the mother. The pregnancy was a full-term without postpartum complications. The baby was born without congenital malformations and with normal neurological status. Undoubtedly, the use of immunosuppressants in pregnancy requires careful planning and monitoring in order to minimize risks and enable favorable outcome.

Conclusion

Pregnancies of kidney transplant recipients are high-risk and require a multidisciplinary approach. Careful clinical follow-up is a prerequisite for a favorable outcome.

Acknowledgement

This work was supported by the Ministry of Education, Science and Technological Development of Serbia (Grant No. 175023).

REFERENCES

- Margoles HR, Gomez-Lobo V, Veis JH, Sherman MJ, Moore J. Successful maternal and fetal outcome in a kidney transplant patient with everolimus exposure throughout pregnancy: A case report. *Transplant Proc* 2014; 46(1): 281–3.
- McKay DB, Josephson MA. Pregnancy after Kidney Transplantation. *Clin J Am Soc Nephrol* 2008; 3(2): 117–25.
- Debska-Slizien A, Galgowska J, Chamienia A, Bulko-Piontecka B, Król E, Lichodziejewska-Niemierko M, et al. Pregnancy after kidney transplantation: a single-center experience and review of the literature. *Transplant Proc* 2014; 46(8): 2668–72.
- NKF KDOQI Guidelines. KDOQI Clinical Practice Guidelines on Hypertension and Antihypertensive Agents in Chronic Kidney Disease. 2015. [cited 2015 May 28]. Available from: http://www2.kidney.org/professionals/KDOQI/guidelines_bp
- Gorgulu N, Yelken B, Caliskan Y, Turkmen A, Sever MS. Does pregnancy increase graft loss in female renal allograft recipients. *Clin Exp Nephrol* 2010; 14(3): 244–7.
- Hodžić E, Brčić M, Kapidžić M, Halilović-Terzić A, Jusufović S, Jasarević A, et al. Pregnancy in renal transplantation. *Med Arch* 2013; 67(3): 215–8.
- Veroux M, Corona D, Veroux P. Pregnancy under everolimus-based Pregnancy under everolimus-based immunosuppression. *Transpl Int* 2011; 24(12): e115–7.
- Pisoni CN, D'Cruz DP. The safety of mycophenolate mofetil in pregnancy. *Expert Opin Drug Saf* 2008; 7(3): 219–22.
- Sijfontis NM, Coscia LA, Constantinescu S, Lavelanet AF, Moritz MJ, Armenti VT. Pregnancy outcomes in solid organ transplant recipients with exposure to mycophenolate mofetil or sirolimus. *Transplantation* 2006; 82(12): 1698–702.

10. Krämer BK, Montagnino G, Del Castillo D, Margreiter R, Sperschneider H, Olbricht CJ, et al. Efficacy and safety of tacrolimus compared with cyclosporin A microemulsion in renal transplantation: 2 year follow-up results. *Nephrol Dial Transplant* 2005; 20(5): 968–73.
11. Surti B, Tan J, Saab S. Pregnancy and liver transplantation. *Liver Int* 2008; 28(9): 1200–6.
12. Flechner SM, Katz AR, Rogers AJ, Van Buren C, Kaban BD. The presence of cyclosporine in body tissues and fluids during pregnancy. *Am J Kidney Dis* 1985; 5(1): 60–3.
13. Hebert MF, Zbeng S, Hays K, Shen DD, Davis CL, Umans JG, et al. Interpreting tacrolimus concentrations during pregnancy and postpartum. *Transplantation* 2013; 95(7): 908–15.
14. Kainz A, Harabacz I, Cowbrick IS, Gadgil S, Haginara D. Analysis of 100 pregnancy outcomes in women treated systemically with tacrolimus. *Transpl Int* 2000; 13 Suppl 1: S299–300.
15. Coscia LA, Constantinescu S, Moritz MJ, Frank AM, Ramirez CB, Maley WR, et al. Report from the National Transplantation Pregnancy Registry (NTPR): Outcomes of pregnancy after transplantation. *Clin Transpl* 2010: 65–85.
16. Briggs GG, Freeman RK, Yaffe SJ. *Drugs in pregnancy and lactation: a reference guide to fetal and neonatal risk*. Philadelphia: Lippincott Williams and Wilkins; 2005. p. 405–7.
17. FDA pregnancy categories. Available from: <http://www.drugs-com/pregnancy-categories.html>
18. Hou S. Pregnancy in renal transplant recipients. *Adv Chronic Kidney Dis* 2013; 20(3): 253–9.
19. van Rummard Heimel PJ, Schobben AF, Huisjes AJ, Franx A, Bruinse HW. The transplacental passage of prednisolone in pregnancies complicated by early-onset HELLP syndrome. *Placenta* 2005; 26(10): 842–5.
20. Poljka JE, Friedman JM. Teratogen update: Azathioprine and 6-mercaptopurine. *Teratology* 2002; 65(5): 240–61.
21. FDA Drug Safety Communication: Risk of oral clefts in children born to mothers taking Topamax (topiramate) Available from: <http://www.fda.gov/Drugs/DrugSafety/ucm245085.htm>
22. Williams D, Mayabi L. Maternal medicines in the fetus. In: *Rodeck CH, Whittle MJ*, editors. *Fetal Medicine: Basic Science and Clinical Practice*. 2nd. London: Churchill Livingstone Elsevier; 2009. p. 167–8.
23. Coté CJ, Meuwissen HJ, Pickering RJ. Effects on the neonate of prednisone and azathioprine administered to the mother during pregnancy. *J Pediatr* 1974; 85(3): 324–8.
24. Goldstein LH, Dolinsky G, Greenberg R, Schaefer C, Cohen-Kerem R, Diar-Citrin O, et al. Pregnancy outcome of women exposed to azathioprine during pregnancy. *Birth Defects Res Part A Clin Mol Teratol* 2007; 79(10): 696–701.
25. Vaziri ND, Ismail M, Martin DC, Gonzales E. Blood coagulation, fibrinolytic and inhibitory profiles in renal transplant recipients: Comparison of cyclosporine and azathioprine. *Int J Artif Organs* 1992; 15(6): 365–9.
26. Vazquez SR, Rondina MT, Pendleton RC. Azathioprine-induced warfarin resistance. *Ann Pharmacother* 2008; 42(7): 1118–23.
27. Armenti VT, Radomski JS, Moritz MJ, Gangban WJ, Hecker WP, Lavelanet A, et al. Report from the National Transplantation Pregnancy Registry (NTPR): Outcomes of pregnancy after transplantation. *Clin Transpl* 2004: 103–14.

Received on December 8, 2015.

Accepted on February 25, 2016.

Online First July, 2016.